

# Maximum Entropy Formulation of the Kirkwood Superposition Approximation

A. Singer\*, Z. Schuss†

April 16, 2004

## Abstract

Using a variational formulation, we derive the Kirkwood superposition approximation for entire space systems at equilibrium. We define the entropy of particle triplets, and show that the Kirkwood closure brings the entropy to its maximal value. The principle of maximum entropy is a well known principle in statistical mechanics and beyond. Our approach leads to a different interpretation for the Kirkwood closure relation, usually explained by probabilistic consideration of dependency and independency of particles. We find the generalization of the Kirkwood closure for finite volume systems with certain boundary conditions as needed to specify any electrochemical or biophysical device, e.g., a concentration cell, battery, or ionic channel. The generalized closure is slightly different than the Kirkwood closure, and enables us to calculate the pair function near the boundary walls of the domain.

## 1 Introduction

The pair correlation function is one of the cornerstones in the theory of simple liquids [1, 3, 2, 4]. Many thermodynamic properties of the fluid can be

---

\*Department of Applied Mathematics, Tel-Aviv University, Ramat-Aviv, Tel-Aviv 69978, Israel, e-mail: amit\_sin@zahav.net.il, partially supported by research grants from the Israel Science Foundation, US-Israel Binational Science Foundation, and DARPA

†Department of Applied Mathematics, Tel-Aviv University, Ramat-Aviv, Tel-Aviv 69978, Israel, e-mail: schuss@post.tau.ac.il, partially supported by research grants from the Israel Science Foundation, US-Israel Binational Science Foundation, and DARPA

derived from the pair function. There are mainly two approaches for finding the pair function. The first approach is based on the Ornstein-Zernike integral equation and a closure relation for the direct and indirect correlation functions. Many closure relations fall into this category, such as the Percus-Yevick approximation (PY), the hypernetted chain approximation (HNC) and the mean spherical approximation (MSA). The second approach relies on the Bogoliubov-Born-Green-Kirkwood-Yvon (BBGKY) hierarchy, which relates the  $n$ -th correlation function with the  $n + 1$ -th correlation function, and assumes a closure relation that connects them. The Kirkwood superposition that approximates the triplet correlation function by the product of the three pair correlation functions, was the first suggested closure of this kind.

Both approaches and the underlying closure relations have their own advantages and disadvantages. Their success is often compared to either molecular dynamics (MD) or Monte-Carlo (MC) simulations. All closures succeed in the limit of (very) low density. However, when the density is increased they tend to fail, sooner or later. Choosing the “best” closure relation is an art in itself. Obviously, each choice of closure relation results in a different approximate solution for the pair function. The BBGKY equation (with or without the superposition approximation) has a great advantage over any other approach. For hard spheres it predicts a point where  $\frac{dp}{d\rho} = 0$ , and hence solidification. Neither the PY or HNC theories can do this.

From a mathematical point of view, all approaches suffer from their lack of error estimates for the approximations being made. This issue is more serious than it may seem. Mathematically, there are no shortage of expansions that do not converge, or converge only after many many terms, or converge in some senses (uniformly, conditionally, asymptotically) and not others. If a few terms of such series are used, and error estimates are not available, it is easy to develop approximations with **qualitative** properties very different from the original function. Scientifically, mathematical issues can sometimes be sidestepped, if the process of scientific investigation (‘guess and check’) converges to an agreed solution. If error estimates are not available, the scientific process is much more difficult and may itself (socially) diverge. Our goal is to develop a mathematically defined closure approximation that has general meaning and allows error estimates.

In this paper we derive the Kirkwood superposition approximation from a variational (Euler-Lagrange) formulation. We define an entropy functional

for the triplets function, and show that it produces the Kirkwood closure (for entire space systems at equilibrium) when it reaches its maximum value, under the constraint that its marginal is the pair function. We hope that this kind of formulation will enable us to find error estimates for this simple closure.

This maximum entropy formulation leads to a different closure for a finite volume system. We write the resulting closure as an integral equation. In the limit of entire space, the solution of this integral equation reduces to the Kirkwood superposition approximation. We describe an iterative procedure for solving the BBGKY equation with this generalized closure.

Using this generalized closure instead of the original Kirkwood closure produces different pair correlation functions. The generalized closure can be used to calculate the pair function in confined geometries. Specifically, we can deduce the resulting pair function near the domain walls (boundaries), and compare it to MC or MD simulations, and to the results of the original Kirkwood closure.

## 2 Maximum Entropy

Let  $\Omega \subset \mathbb{R}^3$ . Suppose  $p_2(\mathbf{x}_1, \mathbf{x}_2)$  is a known symmetric probability distribution function (pdf)

$$\int_{\Omega \times \Omega} p_2(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2 = 1$$

$$p_2(\mathbf{x}_1, \mathbf{x}_2) = p_2(\mathbf{x}_2, \mathbf{x}_1)$$

$$p_2(\mathbf{x}_1, \mathbf{x}_2) \geq 0.$$

In terms of simple liquids statistical mechanics  $p_2(\mathbf{x}_1, \mathbf{x}_2)$  represents the joint pdf of finding two particles at locations  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

Consider the following optimization problem. Find a pdf  $p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$

of the triplets, that satisfies the constraints

$$\begin{aligned}\phi_1(p_3) &= \int_{\Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3 - p_2(\mathbf{x}_1, \mathbf{x}_2) = 0 \\ \phi_2(p_3) &= \int_{\Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_2 - p_2(\mathbf{x}_1, \mathbf{x}_3) = 0 \\ \phi_3(p_3) &= \int_{\Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_1 - p_2(\mathbf{x}_2, \mathbf{x}_3) = 0\end{aligned}$$

which means that  $p_2$  is the marginal of  $p_3$ , by the definition of the marginal probability density, and brings the entropy functional

$$H(p_3) = - \int_{\Omega \times \Omega \times \Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) \log p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \quad (1)$$

to its maximum. This is a variational formulation of the closure problem. To find its solution we use the Lagrange multipliers method. Let  $\lambda_1(\mathbf{x}_2, \mathbf{x}_3)$ ,  $\lambda_2(\mathbf{x}_1, \mathbf{x}_3)$ ,  $\lambda_3(\mathbf{x}_1, \mathbf{x}_2)$  be the multipliers of the constraints, and the functional

$$\begin{aligned}F(p_3, \lambda_1, \lambda_2, \lambda_3) &= H + \lambda_1 \phi_1 + \lambda_2 \phi_2 + \lambda_3 \phi_3 \\ &= - \int_{\Omega \times \Omega \times \Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) \log p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \\ &\quad + \lambda_1(\mathbf{x}_2, \mathbf{x}_3) \left( \int_{\Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_1 - p_2(\mathbf{x}_2, \mathbf{x}_3) \right) \\ &\quad + \lambda_2(\mathbf{x}_1, \mathbf{x}_3) \left( \int_{\Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_2 - p_2(\mathbf{x}_1, \mathbf{x}_3) \right) \\ &\quad + \lambda_3(\mathbf{x}_1, \mathbf{x}_2) \left( \int_{\Omega} p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3 - p_2(\mathbf{x}_1, \mathbf{x}_2) \right).\end{aligned}$$

Since  $F$  is independent of the derivatives of  $p_3$ , the Euler-Lagrange equations

read

$$\nabla_{\mathbf{x}_1} [-\log p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) - 1 + \lambda_1(\mathbf{x}_2, \mathbf{x}_3) + \lambda_2(\mathbf{x}_1, \mathbf{x}_3) + \lambda_3(\mathbf{x}_1, \mathbf{x}_2)] = 0,$$

$$\nabla_{\mathbf{x}_2} [-\log p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) - 1 + \lambda_1(\mathbf{x}_2, \mathbf{x}_3) + \lambda_2(\mathbf{x}_1, \mathbf{x}_3) + \lambda_3(\mathbf{x}_1, \mathbf{x}_2)] = 0,$$

$$\nabla_{\mathbf{x}_3} [-\log p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) - 1 + \lambda_1(\mathbf{x}_2, \mathbf{x}_3) + \lambda_2(\mathbf{x}_1, \mathbf{x}_3) + \lambda_3(\mathbf{x}_1, \mathbf{x}_2)] = 0.$$

Therefore

$$-\log p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) - 1 + \lambda_1(\mathbf{x}_2, \mathbf{x}_3) + \lambda_2(\mathbf{x}_1, \mathbf{x}_3) + \lambda_3(\mathbf{x}_1, \mathbf{x}_2) = C, \quad (2)$$

where  $C$  is a constant, or equivalently

$$p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \gamma_1(\mathbf{x}_2, \mathbf{x}_3)\gamma_2(\mathbf{x}_1, \mathbf{x}_3)\gamma_3(\mathbf{x}_1, \mathbf{x}_2), \quad (3)$$

where

$$\gamma_1(\mathbf{x}_2, \mathbf{x}_3) = e^{\lambda_1(\mathbf{x}_2, \mathbf{x}_3) - (C+1)/3}$$

$$\gamma_2(\mathbf{x}_1, \mathbf{x}_3) = e^{\lambda_2(\mathbf{x}_1, \mathbf{x}_3) - (C+1)/3}$$

$$\gamma_3(\mathbf{x}_1, \mathbf{x}_2) = e^{\lambda_3(\mathbf{x}_1, \mathbf{x}_2) - (C+1)/3}$$

Clearly,  $\gamma_i \geq 0$  ( $i = 1, 2, 3$ ), and therefore  $p_3 \geq 0$ . Moreover,  $\gamma_1 = \gamma_2 = \gamma_3$  because  $p_2$  is symmetric. Let  $\gamma = \gamma_1 = \gamma_2 = \gamma_3$ . Then  $p_3$  takes the form

$$p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \gamma(\mathbf{x}_1, \mathbf{x}_2)\gamma(\mathbf{x}_2, \mathbf{x}_3)\gamma(\mathbf{x}_1, \mathbf{x}_3). \quad (4)$$

We find  $\gamma$  from the constraint that  $p_2$  is the marginal of  $p_3$

$$p_2(\mathbf{x}_1, \mathbf{x}_2) = \gamma(\mathbf{x}_1, \mathbf{x}_2) \int_{\Omega} \gamma(\mathbf{x}_1, \mathbf{x}_3)\gamma(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3. \quad (5)$$

The symmetry of  $p_2$ , i.e.  $p_2(\mathbf{x}_1, \mathbf{x}_2) = p_2(\mathbf{x}_2, \mathbf{x}_1)$  follows that of  $\gamma$ ,  $\gamma(\mathbf{x}_1, \mathbf{x}_2) = \gamma(\mathbf{x}_2, \mathbf{x}_1)$ .

However, in practice, the pdf  $p_2$  is unknown. Nevertheless, it satisfies the BBGKY equation

$$\begin{aligned} 0 = & \mathbf{f}_{\text{ex}}(\mathbf{x}_1)p_2(\mathbf{x}_1, \mathbf{x}_2) + \mathbf{f}(\mathbf{x}_2, \mathbf{x}_1)p_2(\mathbf{x}_1, \mathbf{x}_2) - k_B T \nabla_{\mathbf{x}_1} p_2(\mathbf{x}_1, \mathbf{x}_2) \\ & + (N-2) \int_{\Omega} \mathbf{f}(\mathbf{x}_3, \mathbf{x}_1)p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3, \end{aligned}$$

where  $\mathbf{f}(\mathbf{x}_2, \mathbf{x}_1)$  is the force interacted on a particle located at  $\mathbf{x}_1$ , by another particle located at  $\mathbf{x}_2$ , and  $\mathbf{f}_{\text{ex}}(\mathbf{x}_1)$  is an external force field acting on a particle located at  $\mathbf{x}_1$ . Substituting the maximum entropy closure (4) into the BBGKY equation (6), together with the integral equation (5) results in a system of integral equations for  $p_2$  and  $\gamma$ ,

$$\begin{aligned}
0 &= \mathbf{f}_{\text{ex}}(\mathbf{x}_1)p_2(\mathbf{x}_1, \mathbf{x}_2) + \mathbf{f}(\mathbf{x}_2, \mathbf{x}_1)p_2(\mathbf{x}_1, \mathbf{x}_2) - k_B T \nabla_{\mathbf{x}_1} p_2(\mathbf{x}_1, \mathbf{x}_2) \\
&\quad + (N-2) \int_{\Omega} \mathbf{f}(\mathbf{x}_3, \mathbf{x}_1) \gamma(\mathbf{x}_1, \mathbf{x}_2) \gamma(\mathbf{x}_2, \mathbf{x}_3) \gamma(\mathbf{x}_1, \mathbf{x}_3) d\mathbf{x}_3 \\
p_2(\mathbf{x}_1, \mathbf{x}_2) &= \gamma(\mathbf{x}_1, \mathbf{x}_2) \int_{\Omega} \gamma(\mathbf{x}_1, \mathbf{x}_3) \gamma(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3
\end{aligned} \tag{6}$$

Solving the system (6) will produce the pair function  $p_2$ .

### 3 Two Examples

In this section we discuss two examples in which the system (6) can be solved or simplified.

#### 3.1 Non interacting particles in an external field

Non interacting particles in an external field are described by  $\mathbf{f} \equiv 0$ . Therefore, the system (6) is reduced to

$$0 = \mathbf{f}_{\text{ex}}(\mathbf{x}_1)p_2(\mathbf{x}_1, \mathbf{x}_2) - k_B T \nabla_{\mathbf{x}_1} p_2(\mathbf{x}_1, \mathbf{x}_2) \tag{7}$$

$$p_2(\mathbf{x}_1, \mathbf{x}_2) = \gamma(\mathbf{x}_1, \mathbf{x}_2) \int_{\Omega} \gamma(\mathbf{x}_1, \mathbf{x}_3) \gamma(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3 \tag{8}$$

Solving equation (7) yields

$$p_2(\mathbf{x}_1, \mathbf{x}_2) = e^{-U_{\text{ex}}(\mathbf{x}_1)/k_B T} h(\mathbf{x}_2), \tag{9}$$

where  $\mathbf{f}_{\text{ex}}(\mathbf{x}_1) = -\nabla_{\mathbf{x}_1} U_{\text{ex}}(\mathbf{x}_1)$ , and  $h(\mathbf{x}_2)$  is an arbitrary function (the integration constant). Since  $p_2(\mathbf{x}_1, \mathbf{x}_2) = p_2(\mathbf{x}_2, \mathbf{x}_1)$ , we have

$$p_2(\mathbf{x}_1, \mathbf{x}_2) = C^{-1} e^{-[U_{\text{ex}}(\mathbf{x}_1) + U_{\text{ex}}(\mathbf{x}_2)]/k_B T}, \tag{10}$$

where  $C = \left[ \int_{\Omega} e^{-U \text{ex}(\mathbf{x})/k_B T} d\mathbf{x} \right]^2$  is a normalization constant. As expected, we find that  $\mathbf{x}_1, \mathbf{x}_2$  are independent random variables,  $p_2(\mathbf{x}_1, \mathbf{x}_2) = p_1(\mathbf{x}_1)p_1(\mathbf{x}_2)$ . In this case, the solution to (8) is given by

$$\gamma(\mathbf{x}_1, \mathbf{x}_2) = \sqrt{p_2(\mathbf{x}_1, \mathbf{x}_2)} = \sqrt{p_1(\mathbf{x}_1)}\sqrt{p_1(\mathbf{x}_2)}. \quad (11)$$

Indeed,

$$\begin{aligned} \gamma(\mathbf{x}_1, \mathbf{x}_2) & \int_{\Omega} \gamma(\mathbf{x}_1, \mathbf{x}_3)\gamma(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3 = \\ & = \sqrt{p_1(\mathbf{x}_1)p_1(\mathbf{x}_2)} \int_{\Omega} \sqrt{p_1(\mathbf{x}_1)p_1(\mathbf{x}_3)}\sqrt{p_1(\mathbf{x}_2)p_1(\mathbf{x}_3)} d\mathbf{x}_3 \\ & = p_1(\mathbf{x}_1)p_1(\mathbf{x}_2) \int_{\Omega} p_1(\mathbf{x}_3) d\mathbf{x}_3 = p_1(\mathbf{x}_1, \mathbf{x}_2). \end{aligned}$$

### 3.2 Entire Space Systems

In the limit process  $\Omega \rightarrow \mathbb{R}^3$  all the pdfs tend to zero, so it is more convenient to work with densities. First, let  $\Omega \subset \mathbb{R}^3$  be a bounded domain. The previous example of non-interacting particles give rise to define

$$\delta(\mathbf{x}_1, \mathbf{x}_2) = \frac{\gamma(\mathbf{x}_1, \mathbf{x}_2)}{\sqrt{p_1(\mathbf{x}_1)p_1(\mathbf{x}_2)}}, \quad (12)$$

so equation (8) reads

$$\frac{p_2(\mathbf{x}_1, \mathbf{x}_2)}{p_1(\mathbf{x}_1)p_1(\mathbf{x}_2)} = \delta(\mathbf{x}_1, \mathbf{x}_2) \int_{\Omega} p_1(\mathbf{x}_3)\delta(\mathbf{x}_1, \mathbf{x}_3)\delta(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3. \quad (13)$$

We rewrite equation (13) as

$$\begin{aligned} \frac{p_2^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2)}{p_1^{(\Omega)}(\mathbf{x}_1)p_1^{(\Omega)}(\mathbf{x}_2)} & = \\ \delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2) & \int_{\Omega} p_1^{(\Omega)}(\mathbf{x}_3)\delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_3)\delta^{(\Omega)}(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3, \end{aligned} \quad (14)$$

where  $p_2^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2) = p_2(\mathbf{x}_1, \mathbf{x}_2)$ ,  $\delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2) = \delta(\mathbf{x}_1, \mathbf{x}_2)$ ,  $p_1^{(\Omega)}(\mathbf{x}_1) = p_1(\mathbf{x}_1)$ , to emphasize their dependency on the specific domain  $\Omega$ . Let

$$g_2(\mathbf{x}_1, \mathbf{x}_2) = \lim_{\Omega \rightarrow \mathbb{R}^3} \frac{p_2^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2)}{p_1^{(\Omega)}(\mathbf{x}_1)p_1^{(\Omega)}(\mathbf{x}_2)}. \quad (15)$$

For example, if the two particles become independent when they are separated,

$$p_2^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2) = p_1^{(\Omega)}(\mathbf{x}_1)p_1^{(\Omega)}(\mathbf{x}_2)(1 + o(1)), \text{ for } |\mathbf{x}_1 - \mathbf{x}_2| \gg 1, \quad (16)$$

then  $\lim_{|\mathbf{x}_2| \rightarrow \infty} g_2(\mathbf{x}_1, \mathbf{x}_2) = 1$ .

Next, we show that  $\lim_{\Omega \rightarrow \mathbb{R}^3} \delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2) = g_2(\mathbf{x}_1, \mathbf{x}_2)$ , given the assumption (16). Indeed,

$$\begin{aligned} \int_{\Omega} p_1(\mathbf{x}_3) \frac{p_2(\mathbf{x}_1, \mathbf{x}_3)}{p_1(\mathbf{x}_1)p_1(\mathbf{x}_3)} \frac{p_2(\mathbf{x}_2, \mathbf{x}_3)}{p_1(\mathbf{x}_2)p_1(\mathbf{x}_3)} d\mathbf{x}_3 &= \int_{\Omega} p_1(\mathbf{x}_3) (1 + o(1)) d\mathbf{x}_3 \\ &= 1 + o(1). \end{aligned} \quad (17)$$

Taking the limit  $\Omega \rightarrow \mathbb{R}^3$  the  $o(1)$  term vanishes, and equation (14) follows

$$\delta(\mathbf{x}_1, \mathbf{x}_2) = g_2(\mathbf{x}_1, \mathbf{x}_2), \quad (18)$$

as asserted.

We interpret equation (18) as the Kirkwood superposition approximation. Equations (4) and (12) implies that the triplet pdf satisfies

$$\frac{p_3^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)}{p_1^{(\Omega)}(\mathbf{x}_1)p_1^{(\Omega)}(\mathbf{x}_2)p_1^{(\Omega)}(\mathbf{x}_3)} = \delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2)\delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_3)\delta^{(\Omega)}(\mathbf{x}_2, \mathbf{x}_3). \quad (19)$$

Taking the limit  $\Omega \rightarrow \mathbb{R}^3$ , using equation (18), we obtain

$$\begin{aligned} g_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) &= \lim_{\Omega \rightarrow \mathbb{R}^3} \frac{p_3^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)}{p_1^{(\Omega)}(\mathbf{x}_1)p_1^{(\Omega)}(\mathbf{x}_2)p_1^{(\Omega)}(\mathbf{x}_3)} \\ &= g_2(\mathbf{x}_1, \mathbf{x}_2)g_2(\mathbf{x}_1, \mathbf{x}_3)g_2(\mathbf{x}_2, \mathbf{x}_3), \end{aligned} \quad (20)$$

which is the Kirkwood closure relation. Usually, the motivation for using the Kirkwood superposition approximation is a probabilistic consideration of dependency and independency of particles (see also Section 5.) Here we find another interpretation for the Kirkwood closure.

The Kirkwood closure is the (only) closure relation that brings the entropy of triplets of particles to its maximum value. The principle of maximum entropy is a well known principle in statistical mechanics, in testing statistical hypotheses [5, 6] and beyond. In the following section we give further motivation for its use.



## 4 Minimum Helmholtz Free Energy

In the beginning of every elementary textbook in statistical mechanics, it is mentioned that the Boltzmann distribution

$$p_N(\mathbf{x}_1, \dots, \mathbf{x}_N) = \frac{1}{Z_N} e^{-U(\mathbf{x}_1, \dots, \mathbf{x}_N)/k_B T}, \quad (21)$$

brings the Helmholtz free energy

$$\begin{aligned} F(p) &= U(p) - k_B T H(p) \\ &= \int_{\Omega^N} U(\mathbf{x}_1, \dots, \mathbf{x}_N) p(\mathbf{x}_1, \dots, \mathbf{x}_N) d\mathbf{x}_1 \cdots d\mathbf{x}_N \\ &\quad + k_B T \int_{\Omega^N} p(\mathbf{x}_1, \dots, \mathbf{x}_N) \log p(\mathbf{x}_1, \dots, \mathbf{x}_N) d\mathbf{x}_1 \cdots d\mathbf{x}_N, \end{aligned} \quad (22)$$

to its minimum under the normalization constraint

$$\int_{\Omega^N} p(\mathbf{x}_1, \dots, \mathbf{x}_N) d\mathbf{x}_1 \cdots d\mathbf{x}_N = 1. \quad (23)$$

For a pairwise additive potential together with an external field force

$$U(\mathbf{x}_1, \dots, \mathbf{x}_N) = \sum_{1 \leq i < j \leq N} U(\mathbf{x}_i, \mathbf{x}_j) + \sum_{j=1}^N U_{\text{ex}}(\mathbf{x}_j), \quad (24)$$

the potential energy term  $U(p)$  of the Helmholtz free energy (22), takes the simple form

$$\begin{aligned} U(p) &= \int_{\Omega^N} \left( \sum_{1 \leq i < j \leq N} U(\mathbf{x}_i, \mathbf{x}_j) + \sum_{j=1}^N U_{\text{ex}}(\mathbf{x}_j) \right) p(\mathbf{x}_1, \dots, \mathbf{x}_N) d\mathbf{x}_1 \cdots d\mathbf{x}_N \\ &= \frac{N(N-1)}{2} \int_{\Omega^2} U(\mathbf{x}_1, \mathbf{x}_2) p_2(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2 + N \int_{\Omega} U_{\text{ex}}(\mathbf{x}_1) p_1(\mathbf{x}_1) d\mathbf{x}_1, \end{aligned}$$

where

$$\begin{aligned} p_2(\mathbf{x}_1, \mathbf{x}_2) &= \int_{\Omega^{N-2}} p_N(\mathbf{x}_1, \dots, \mathbf{x}_N) d\mathbf{x}_1 \cdots d\mathbf{x}_N, \\ p_1(\mathbf{x}_1) &= \int_{\Omega} p_2(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_2, \end{aligned}$$

are the marginal distributions. If the pdf  $p_2(\mathbf{x}_1, \mathbf{x}_2)$  is assumed to be known, as in Section 2, then the energy term of the Helmholtz free energy  $U(p)$  is also known. Therefore, minimizing the Helmholtz free energy, under the assumption that the pdf  $p_2(\mathbf{x}_1, \mathbf{x}_2)$  is known, is equivalent for maximizing the entropy, since  $U(p)$  is constant.

## 5 Probabilistic Interpretation of the Kirkwood Closure

The Kirkwood superposition approximation (20) was the first closure relation to be suggested and tested in the theory of simple liquids. This fact might be explained by its simplicity and its intuitive origin. In this section we give a probabilistic interpretation of the Kirkwood superposition approximation, and find its generalization for higher levels closure relations of the BBGKY hierarchy. The problem at level  $n$  ( $n \geq 2$ ) is to find an approximation for the  $n + 1$ -particle pdf in terms of the  $n$ -particle pdf. For example, the Kirkwood superposition approximation (20) closes the hierarchy at level  $n = 2$ . High orders closures are expected to be much more accurate, fitting better the experimental and simulated (MC or MD) data **REFERENCES**. However, the computational complexity increases drastically with  $n$ .

First, consider the case  $n = 2$ . We assume that particles become independent as they are distanced (16). In order to make the exact equality (16) into an approximation, we assume that there exists a distance  $d > 0$  such that

$$p_2(\mathbf{x}_1, \mathbf{x}_2) = p_1(\mathbf{x}_1)p_1(\mathbf{x}_2), \quad (25)$$

for  $|\mathbf{x}_1 - \mathbf{x}_2| > d$ . Three interchangeable particles can be in four different configurations with respect to the distance  $d$ , depending on the number of intersections (see Figure 1). In all configurations but configuration (d), where all three particles intersect, there are at least two particles that do not intersect. Since the particles are interchangeable we may assume that  $|\mathbf{x}_1 - \mathbf{x}_3| > d$ . Applying Bayes' law we have

$$p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = p_3(\mathbf{x}_3|\mathbf{x}_1, \mathbf{x}_2)p_2(\mathbf{x}_1, \mathbf{x}_2). \quad (26)$$

By the independency assumption (25) we have

$$p_3(\mathbf{x}_3|\mathbf{x}_1, \mathbf{x}_2) = p_2(\mathbf{x}_3|\mathbf{x}_2), \quad (27)$$




Figure 1: Four configurations of three particles. (a) no intersections (b) one intersection (c) two intersections (d) three intersections.

therefore,

$$p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = p_2(\mathbf{x}_3|\mathbf{x}_2)p_2(\mathbf{x}_1, \mathbf{x}_2) = \frac{p_2(\mathbf{x}_2, \mathbf{x}_3)}{p_1(\mathbf{x}_2)}p_2(\mathbf{x}_1, \mathbf{x}_2). \quad (28)$$

Multiplying by  $1 = \frac{p_2(\mathbf{x}_1, \mathbf{x}_3)}{p_1(\mathbf{x}_1)p_1(\mathbf{x}_3)}$  we obtain

$$p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \frac{p_2(\mathbf{x}_1, \mathbf{x}_2)p_2(\mathbf{x}_2, \mathbf{x}_3)p_2(\mathbf{x}_1, \mathbf{x}_3)}{p_1(\mathbf{x}_1)p_1(\mathbf{x}_2)p_1(\mathbf{x}_3)}, \quad (29)$$

which is the Kirkwood superposition approximation. We see that the Kirkwood closure is a good approximation when at least two particles are distant. However, it fails when all three particles are close to each other.

Next, we find the  $n$ -level Kirkwood closure relation for the  $n + 1$ -particle pdf in terms of the  $n$ -particle pdf, using probabilistic considerations.

**Proposition:** *The  $n$ -level Kirkwood closure relation is given by*

$$p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} p_k(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}}. \quad (30)$$

**Proof:** We have already seen that the approximation holds for  $n = 2, 3$ . Assuming, by induction, that at least two particles are distant, we can assume without loss of generality, that particles 1 and  $n$  are far apart,  $|\mathbf{x}_1 - \mathbf{x}_n| > d$ . Using Bayes' law, we find that

$$p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = p_{n-1}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n-1})p_n(\mathbf{x}_n|\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n-1}).$$

Since particles 1 and  $n$  are far apart it follows that

$$p_n(\mathbf{x}_n|\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n-1}) = p_{n-1}(\mathbf{x}_n|\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{n-1}) = \frac{p_{n-1}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_n)}{p_{n-2}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{n-1})}.$$

Hence,

$$p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \frac{p_{n-1}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n-1})p_{n-1}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_n)}{p_{n-2}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{n-1})}. \quad (31)$$

It follows from the induction assumption that for every  $j = 2, 3, \dots, n-1$ , with particles 1 and  $n$  far apart, we have

$$\begin{aligned} 1 &= p_{n-1}(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_n) \\ &\times \prod_{k=1}^{n-2} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n, i_i \neq j} p_k(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}}. \end{aligned} \quad (32)$$

Multiplying eqs.(31) and (32) (for all  $j = 2, 3, \dots, n-1$ ) ends the proof.  $\diamond$

**Corollary:** For  $n = 4$  Kirkwood's formula becomes

$$\begin{aligned} &p_4(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4) \\ &= \frac{p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_4)p_3(\mathbf{x}_1, \mathbf{x}_3, \mathbf{x}_4)p_3(\mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4)p_1(\mathbf{x}_1)p_1(\mathbf{x}_2)p_1(\mathbf{x}_3)p_1(\mathbf{x}_4)}{p_2(\mathbf{x}_1, \mathbf{x}_2)p_2(\mathbf{x}_1, \mathbf{x}_3)p_2(\mathbf{x}_1, \mathbf{x}_4)p_2(\mathbf{x}_2, \mathbf{x}_3)p_2(\mathbf{x}_2, \mathbf{x}_4)p_2(\mathbf{x}_3, \mathbf{x}_4)}. \end{aligned} \quad (33)$$

In the case of the entire space  $\Omega = \mathbb{R}^3$ , we define

$$g_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \lim_{\Omega \rightarrow \mathbb{R}^3} \frac{p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)}{\prod_{j=1}^n p_1(\mathbf{x}_j)}. \quad (34)$$

Dividing equation (30) by  $\prod_{j=1}^n p_1(\mathbf{x}_j)$  gives

$$\frac{p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)}{\prod_{j=1}^n p_1(\mathbf{x}_j)} = \prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \left( \frac{p_k(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})}{\prod_{j=1}^k p_1(\mathbf{x}_{i_j})} \right)^{(-1)^{n-1-k}}, \quad (35)$$

where we used the combinatorial identity

$$\sum_{k=1}^{n-1} \binom{n-1}{k-1} (-1)^{n-1-k} = 1. \quad (36)$$

Note that the  $k = 1$  terms in the product of equation (35) cancel out, so the product may begin from  $k = 2$

$$\frac{p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)}{\prod_{j=1}^n p_1(\mathbf{x}_j)} = \prod_{k=2}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \left( \frac{p_k(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})}{\prod_{j=1}^k p_1(\mathbf{x}_{i_j})} \right)^{(-1)^{n-1-k}}. \quad (37)$$

Taking the limit  $\Omega \rightarrow \mathbb{R}^3$  we obtain the  $n$ -level Kirkwood closure relation

$$g_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \prod_{k=2}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} g_k(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}}. \quad (38)$$

Examples:

- $n = 3$

$$g_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = g_2(\mathbf{x}_1, \mathbf{x}_2)g_2(\mathbf{x}_1, \mathbf{x}_3)g_2(\mathbf{x}_2, \mathbf{x}_3)$$

- $n = 4$

$$g_4(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4) = \frac{g_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)g_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_4)g_3(\mathbf{x}_1, \mathbf{x}_3, \mathbf{x}_4)g_3(\mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4)}{g_2(\mathbf{x}_1, \mathbf{x}_2)g_2(\mathbf{x}_1, \mathbf{x}_3)g_2(\mathbf{x}_1, \mathbf{x}_4)g_2(\mathbf{x}_2, \mathbf{x}_3)g_2(\mathbf{x}_2, \mathbf{x}_4)g_2(\mathbf{x}_3, \mathbf{x}_4)}$$

## 6 High Level Entropy Closure

In this section we use the maximum entropy principle to derive the  $n$ -level closure relation, and compare the resulting closure relation with the  $n$ -level probabilistic Kirkwood closure of Section 5. The problem at level  $n$  ( $n \geq 2$ ) is to find an approximation for the  $n + 1$ -particle pdf in terms of the  $n$ -particle pdf. For example, the Kirkwood superposition approximation (20) closes the hierarchy at level  $n = 2$ . We use the principle of maximum entropy to obtain the closure relation. Similar to the derivation of Section 2 we assume that the  $n$ -particle pdf  $p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  is known, and we search for the  $n + 1$ -particle pdf  $p_{n+1}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1})$  that maximizes the entropy

$$H = - \int_{\Omega^{n+1}} p_{n+1}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1}) \log p_{n+1}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1}) d\mathbf{x}_1 d\mathbf{x}_2 \cdots d\mathbf{x}_{n+1}, \quad (39)$$

with the  $n + 1$  constraints that  $p_n$  is the marginal of  $p_{n+1}$

$$p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \int_{\Omega} p_{n+1}(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_{n+1}) d\mathbf{x}_j, \quad j = 1, 2, \dots, n + 1. \quad (40)$$

Since  $p_2$  is the marginal of  $p_n$  ( $n \geq 2$ ), it follows that  $p_2$  is also known. Therefore, for a pairwise additive potential, maximizing the Helmholtz free energy of the  $n + 1$ -particle system is equivalent to minimizing the entropy of the  $n + 1$ -particle system. Introducing the Lagrange multipliers  $\lambda_j(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_{n+1})$ ,  $j = 1, 2, \dots, n+1$ , the Euler-Lagrange equation gives

$$-\log p_{n+1} - 1 + \sum_{j=1}^{n+1} \lambda_j = 0. \quad (41)$$

Since the  $n$  particles are interchangeable,

$$p_{n+1}(\mathbf{x}_1, \dots, \mathbf{x}_{n+1}) = \prod_{j=1}^{n+1} \gamma(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_{n+1}). \quad (42)$$

Integration with respect to  $\mathbf{x}_{n+1}$  yields

$$p_n(\mathbf{x}_1, \dots, \mathbf{x}_n) = \gamma(\mathbf{x}_1, \dots, \mathbf{x}_n) \int_{\Omega} \prod_{j=1}^n \gamma(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_{n+1}) d\mathbf{x}_{n+1}. \quad (43)$$

Solving the non-linear integral equation (43) for  $\gamma$  and substituting in equation (42) is the  $n$ -level closure relation of the  $n$ -level BBGKY hierarchy equation for  $p_n$

$$\begin{aligned} 0 &= \mathbf{f}_{\text{ex}}(\mathbf{x}_1) p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) + \sum_{j=2}^n \mathbf{f}(\mathbf{x}_j, \mathbf{x}_1) p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \\ &\quad - k_B T \nabla_{\mathbf{x}_1} p_n(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \\ &\quad + (N - n) \int_{\Omega} \mathbf{f}(\mathbf{x}_3, \mathbf{x}_1) p_{n+1}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1}) d\mathbf{x}_{n+1}. \end{aligned} \quad (44)$$

## 6.1 Entire Space Systems

We have seen in Section 3 that for entire space systems  $\Omega = \mathbb{R}^3$ , the maximum entropy principle yields the Kirkwood superposition approximation ( $n = 2$ ).

In this section we show that the maximum entropy principle results in the probabilistic Kirkwood closure (30) for all levels  $n \geq 2$ .

First, let  $\Omega \subset \mathbb{R}^3$  be a bounded domain. Let

$$h_n^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \frac{p_n^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)}{\prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}}}, \quad (45)$$

and

$$\delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \frac{\gamma^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)}{\prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k} \frac{n-k}{n-k+1}}}. \quad (46)$$

Dividing equation (43) by  $\prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}}$  we obtain

$$\begin{aligned} h_n^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) &= \delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \int_{\Omega} F(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n, \mathbf{x}_{n+1}) \\ &\quad \times \prod_{j=1}^n \delta^{(\Omega)}(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_{n+1}) d\mathbf{x}_{n+1}, \end{aligned} \quad (47)$$

where

$$\begin{aligned} F(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n, \mathbf{x}_{n+1}) &= \prod_{k=1}^{n-2} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}(n-k-1)} \\ &\quad \times \prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_{k-1} \leq n} p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{k-1}}, \mathbf{x}_{n+1})^{(-1)^{n-1-k}(n-k)}. \end{aligned}$$

For  $\min_{1 \leq j \leq n} |\mathbf{x}_{n+1} - \mathbf{x}_j| \gg 1$ , the  $n+1$ -th particle becomes independent of particles  $1, 2, \dots, n$ , and we have

$$p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{k-1}}, \mathbf{x}_{n+1}) = p_1^{(\Omega)}(\mathbf{x}_{n+1}) p_{k-1}^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{k-1}}) (1 + o(1)), \quad (48)$$

for all  $k = 1, 2, \dots, n$  and all sets of indices  $1 \leq i_1 < i_2 < \dots < i_{k-1} \leq n$ .

Therefore, to leading order

$$\begin{aligned}
F(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n, \mathbf{x}_{n+1}) &= \prod_{k=1}^{n-2} \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n} p_k^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-1-k}(n-k-1)} \\
&\times \prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_{k-1} \leq n} p_{k-1}^{(\Omega)}(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{k-1}})^{(-1)^{n-1-k}(n-k)} \\
&\times \prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_{k-1} \leq n} p_1^{(\Omega)}(\mathbf{x}_{n+1})^{(-1)^{n-1-k}(n-k)} \\
&= \prod_{k=1}^{n-1} \prod_{1 \leq i_1 < i_2 < \dots < i_{k-1} \leq n} p_1^{(\Omega)}(\mathbf{x}_{n+1})^{(-1)^{n-1-k}(n-k)} \\
&= p_1^{(\Omega)}(\mathbf{x}_{n+1}) \sum_{k=1}^{n-1} (-1)^{n-1-k}(n-k) \binom{n}{k-1} \\
&= p_1^{(\Omega)}(\mathbf{x}_{n+1}), \tag{49}
\end{aligned}$$

where we have used the combinatorial identity

$$\sum_{k=1}^{n-1} (-1)^{n-1-k}(n-k) \binom{n}{k-1} = 1. \tag{50}$$

We conclude that

$$\begin{aligned}
h_n^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) &= \delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \int_{\Omega} p_1^{(\Omega)}(\mathbf{x}_{n+1}) (1 + o(1)) \\
&\times \prod_{j=1}^n \delta^{(\Omega)}(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_{n+1}) d\mathbf{x}_{n+1}.
\end{aligned}$$

Therefore,

$$\lim_{\Omega \rightarrow \mathbb{R}^3} \delta^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \lim_{\Omega \rightarrow \mathbb{R}^3} h_n^{(\Omega)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \tag{51}$$

Substituting eq. (46) in eq. (42) while using the relations (45) and (51), and the definition (34) we find that

$$g_{n+1}(\mathbf{x}_1, \dots, \mathbf{x}_{n+1}) = \prod_{k=2}^n \prod_{1 \leq i_1 < i_2 < \dots < i_k \leq n+1} g_k(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_k})^{(-1)^{n-k}}. \tag{52}$$

We observe that for entire space systems, the probabilistic Kirkwood closure (38) agrees with the maximum entropy closure (52) for all orders  $n$ .



## 6.2 Closure at the highest level $n = N - 1$

Although the probabilistic Kirkwood closure and the maximum entropy closure agree in the case of entire space systems, they differ in the case of confined systems of finite number of particles  $N$ . It appears that the maximum entropy closure (42) is exact when applying it at the highest level  $n = N - 1$ , while the probabilistic Kirkwood closure (30) is not exact. In other words, the maximum entropy closure relation yields the Boltzmann distribution (21), and what thought at first to be an approximation, turns out to be the exact result. Indeed, let

$$\begin{aligned} & \gamma(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{N-1}) \\ &= \left( \frac{1}{Z_N} \right)^{1/N} \exp \left\{ - \left[ \frac{1}{N-2} \sum_{1 \leq i < j \leq N-1} U(\mathbf{x}_i, \mathbf{x}_j) + \frac{1}{N-1} \sum_{j=1}^{N-1} U_{\text{ex}}(\mathbf{x}_j) \right] / k_B T \right\}. \end{aligned} \quad (53)$$

Clearly,

$$\prod_{j=1}^N \gamma(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_j, \dots, \mathbf{x}_N) = \frac{1}{Z_N} e^{-U(\mathbf{x}_1, \dots, \mathbf{x}_N)/k_B T}, \quad (54)$$

which is the Boltzmann distribution. The Boltzmann distribution obviously satisfies the BBGKY equation (44). Therefore, we have found a solution to BBGKY equation which satisfies the closure relation (42). This solution coincides with the Boltzmann distribution, and so we conclude that it is the exact solution. For the  $N - 1$ -particle pdf  $p_{N-1}$  we have

$$p_{N-1}(\mathbf{x}_1, \dots, \mathbf{x}_{N-1}) = \frac{1}{Z_N} \int_{\Omega} e^{-U(\mathbf{x}_1, \dots, \mathbf{x}_N)/k_B T} d\mathbf{x}_N, \quad (55)$$

which is both the exact and "approximated" result.

The probabilistic Kirkwood approximation (30) gives however a different result. If it were to be exact, then its resulted  $N - 1$ -particle pdf must have been given by equation (55). Therefore, all lower level pdf's  $p_n$  ( $n \leq N - 1$ ) are (multiple) integrals of the Boltzmann distribution. Therefore, by the closure (30),  $p_N$  should have been a multiplication of integrals of the Boltzmann distribution, which is a contradiction to the known form of the Boltzmann distribution (21).

Although the maximum entropy closure is exact at the highest order  $n = N - 1$ , it is not exact at lower orders. The observation made here

motivates us to believe that the maximum entropy closure will turn out to be more accurate (i.e., fit the experimental data better) than the probabilistic Kirkwood closure in confined systems, even when used at lower levels ( $n = 2, 3$ ).

## 7 Confined Systems

Systems in bounded domains are particularly important, because only bounded domains can include the spatially nonuniform boundary conditions needed to describe devices, with spatially distinct inputs, outputs, and (sometimes) power supplies. A large fraction of electrochemistry involves such devices as batteries or concentration cells. A large fraction of molecular biology involves such devices as proteins that transport ions across otherwise impermeable membranes [7, 8].

In the general case, where  $\Omega \subset \mathbb{R}^3$  is a bounded domain, there is no analytic solution to the system (6). We propose to solve this system by the following iterative scheme

1. Initial guess  $\gamma^{(0)}(\mathbf{x}_1, \mathbf{x}_2)$ . Set  $i = 0$ .
2. Solve for  $p_2^{(i)}(\mathbf{x}_1, \mathbf{x}_2)$  the non-homogeneous linear equation

$$\begin{aligned} & k_B T \nabla_{\mathbf{x}_1} p_2^{(i)}(\mathbf{x}_1, \mathbf{x}_2) - \mathbf{f}(\mathbf{x}_2, \mathbf{x}_1) p_2^{(i)}(\mathbf{x}_1, \mathbf{x}_2) - \mathbf{f}_{\text{EX}}(\mathbf{x}_1) p_2^{(i)}(\mathbf{x}_1, \mathbf{x}_2) \\ &= (N - 2) \int_{\Omega} \mathbf{f}(\mathbf{x}_3, \mathbf{x}_1) \gamma^{(i)}(\mathbf{x}_1, \mathbf{x}_2) \gamma^{(i)}(\mathbf{x}_2, \mathbf{x}_3) \gamma^{(i)}(\mathbf{x}_1, \mathbf{x}_3) d\mathbf{x}_3. \end{aligned}$$

3. Solve the non-linear system for  $\gamma^{(i+1)}$

$$p_2^{(i)}(\mathbf{x}_1, \mathbf{x}_2) = \gamma^{(i+1)}(\mathbf{x}_1, \mathbf{x}_2) \int_{\Omega} \gamma^{(i+1)}(\mathbf{x}_2, \mathbf{x}_3) \gamma^{(i+1)}(\mathbf{x}_1, \mathbf{x}_3) d\mathbf{x}_3. \quad (56)$$

4.  $i \leftarrow i + 1$ . Return to step 2, until convergence.

The analysis of the previous section indicates that a good initial guess might be

$$\gamma^{(0)}(\mathbf{x}_1, \mathbf{x}_2) = \sqrt{p_1(\mathbf{x}_1) p_1(\mathbf{x}_2)} g_2(\mathbf{x}_1, \mathbf{x}_2), \quad (57)$$

where  $g_2(\mathbf{x}_1, \mathbf{x}_2)$  is the solution to the BBGKY equation with the Kirkwood superposition approximation in the entire space  $\mathbb{R}^3$ . This solution can be

found rather easily using the inherited symmetries of the problem. For example, it is well known that if  $\mathbf{f}_{\text{EX}} = 0$ , then  $g_2(\mathbf{x}_1, \mathbf{x}_2) = g_2(|\mathbf{x}_1 - \mathbf{x}_2|)$ , and the problem for  $g_2$  becomes one dimensional.

Step 2 requires the solution of a linear partial differential equation in a bounded region. This equation can be written in a gradient form

$$\begin{aligned} \nabla_{\mathbf{x}_1} \left[ e^{(U(\mathbf{x}_1, \mathbf{x}_2) + U_{\text{EX}}(\mathbf{x}_1))/k_B T} p_2(\mathbf{x}_1, \mathbf{x}_2) \right] = \\ - \frac{N-2}{k_B T} e^{(U(\mathbf{x}_1, \mathbf{x}_2) + U_{\text{EX}}(\mathbf{x}_1))/k_B T} \int_{\Omega} \nabla_{\mathbf{x}_1} U(\mathbf{x}_1, \mathbf{x}_3) p_3(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3. \end{aligned}$$

The identity  $\nabla_{\mathbf{x}_1} \times \nabla_{\mathbf{x}_1} u(\mathbf{x}_1) = 0$ , for all  $u$ , imposes a solvability condition for  $\gamma$ . Indeed, taking the curl of the last equation, together with the closure (4) result in

$$\begin{aligned} \nabla_{\mathbf{x}_1} \times \left[ e^{(U(\mathbf{x}_1, \mathbf{x}_2) + U_{\text{EX}}(\mathbf{x}_1))/k_B T} \int_{\Omega} \nabla_{\mathbf{x}_1} U(\mathbf{x}_1, \mathbf{x}_3) \gamma(\mathbf{x}_1, \mathbf{x}_2) \gamma(\mathbf{x}_1, \mathbf{x}_3) \gamma(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3 \right] \\ = 0. \end{aligned}$$

In step 3 we solve a non-linear integral equation. We suggest solving the non-linear equation (5) by a Newton-Raphson iterative scheme. Let  $\gamma^{(n)}(\mathbf{x}_1, \mathbf{x}_2)$  be the  $n$ -th iteration. Define the operator  $\Gamma^{(n)} : \Omega^2 \rightarrow \Omega^2$  as follows

$$\Gamma^{(n)} u(\mathbf{x}, \mathbf{z}) = \int_{\Omega} \gamma^{(n)}(\mathbf{x}, \mathbf{y}) u(\mathbf{z}, \mathbf{y}) d\mathbf{y}. \quad (58)$$

Let the operator  $S : \Omega^2 \rightarrow \Omega^2$  be the symmetrization operator

$$Su(\mathbf{x}, \mathbf{y}) = u(\mathbf{y}, \mathbf{x}). \quad (59)$$

The Newton-Raphson iteration scheme suggests

$$\gamma^{(n+1)} = \gamma^{(n)} + \Delta(\mathbf{x}_1, \mathbf{x}_2), \quad (60)$$

where  $\Delta(\mathbf{x}_1, \mathbf{x}_2)$  satisfies the linear integral equation

$$\begin{aligned} p_2(\mathbf{x}_1, \mathbf{x}_2) - p_2^{(n)}(\mathbf{x}_1, \mathbf{x}_2) &= \frac{p_2^{(n)}(\mathbf{x}_1, \mathbf{x}_2)}{\gamma^{(n)}(\mathbf{x}_1, \mathbf{x}_2)} \Delta(\mathbf{x}_1, \mathbf{x}_2) \\ &+ \gamma^{(n)}(\mathbf{x}_1, \mathbf{x}_2) \Gamma^{(n)} \Delta(\mathbf{x}_1, \mathbf{x}_2) \\ &+ \gamma^{(n)}(\mathbf{x}_1, \mathbf{x}_2) S \Gamma^{(n)} \Delta(\mathbf{x}_1, \mathbf{x}_2), \end{aligned}$$

where

$$p_2^{(n)}(\mathbf{x}_1, \mathbf{x}_2) = \gamma^{(n)}(\mathbf{x}_1, \mathbf{x}_2) \int_{\Omega} \gamma^{(n)}(\mathbf{x}_1, \mathbf{x}_3) \gamma^{(n)}(\mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3. \quad (61)$$

We may write the iteration equivalently as

$$\gamma^{(n+1)} = \gamma^{(n)} + \left( \frac{p_2^{(n)}}{\gamma^{(n)}} + \gamma^{(n)} \Gamma^{(n)} + \gamma^{(n)} S \Gamma^{(n)} \right)^{-1} (p_2 - p_2^{(n)}). \quad (62)$$

The algorithm steps are performed until convergence is achieved.

We have yet to test our generalized Kirkwood closure in practice. The resulting pair correlation function should be compared with MD or MC simulations of particles in a confined region. We can only hope that it will outperform the regular Kirkwood superposition approximation in bounded domains. The observation of subsection 6.2 and the generality of the maximum entropy principle (minimum Helmholtz free energy) motivates this belief. We expect to see the difference between the two closures near the boundary walls of the domain.

## 8 Mixtures

The maximum entropy principle is also applicable to find closure relations of mixtures, both in confined domains and in the entire space. Suppose a mixture of  $S \geq 2$  species, with  $N_\alpha$  ( $\alpha = 1, 2, \dots, S$ ) particles of each specie. Let  $N = \sum_{\alpha=1}^S N_\alpha$  be the total number of particles of all specie. There are  $S^2$  2-particle pdfs,

$$p_2^{\alpha\beta}(\mathbf{x}_1, \mathbf{x}_2) \quad \alpha, \beta \in \{1, 2, \dots, S\},$$

that exhibit the symmetry  $p_2^{\alpha\beta}(\mathbf{x}_1, \mathbf{x}_2) = p_2^{\beta\alpha}(\mathbf{x}_2, \mathbf{x}_1)$ . In this section we briefly discuss how to find the closure relation in the mixture problem.

In the maximum entropy approach, one is searching for  $S^3$  3-particle pdfs  $p_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$ ,  $\alpha, \beta, \gamma \in \{1, 2, \dots, S\}$ , that bring the entropy

$$H = - \sum_{\alpha, \beta, \gamma=0}^S \frac{N_\alpha}{N} \frac{N_\beta}{N} \frac{N_\gamma}{N} \int_{\Omega} p_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) \log p_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \quad (63)$$

to maximum, with the  $3S^3$  marginal constraints

$$\begin{aligned}
p_2^{\alpha\beta}(\mathbf{x}_1, \mathbf{x}_2) &= \int_{\Omega} p_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_3, \\
p_2^{\alpha\gamma}(\mathbf{x}_1, \mathbf{x}_3) &= \int_{\Omega} p_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_2, \\
p_2^{\beta\gamma}(\mathbf{x}_2, \mathbf{x}_3) &= \int_{\Omega} p_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) d\mathbf{x}_1.
\end{aligned} \tag{64}$$

This variational problem is solved using the Euler-Lagrange formulation similar to the derivation done in Section 2.

In the case of an entire space system, it can be shown in a similar way to the method of subsections 3.2 and 6.1, that the mixture entropy closure coincides with the probabilistic Kirkwood closure. For example, they both suggests that the triplets correlation functions are related to the pair correlation function via

$$g_3^{\alpha\beta\gamma}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = g_2^{\alpha\beta}(\mathbf{x}_1, \mathbf{x}_2)g_2^{\alpha\gamma}(\mathbf{x}_1, \mathbf{x}_3)g_2^{\beta\gamma}(\mathbf{x}_2, \mathbf{x}_3), \tag{65}$$

for  $\alpha, \beta, \gamma \in \{1, 2, \dots, S\}$ . Similarly, one can obtain also high order closures.

In confined systems, the Euler-Lagrange formulation leads to integral equation of the form (5). Note that since the entropy (63) depends on the particle fraction  $\frac{N_\alpha}{N}$ , we expect the resulting confined system pair correlation also to depend on the particle fraction.

## 9 Discussion and Summary

We have used the maximum entropy principle to derive a closure relation for the BBGKY hierarchy. It is possible to consider functionals other than the entropy functional that will yield different (known) closures. In fact, a somewhat similar approach is exercised in the density functional theory (DFT). In the DFT problem setup, one assumes a functional of the pair correlation function (e.g., the Helmholtz free energy) to be maximized. The function that brings the given functional to its maximal value is the resulting pair correlation function. Using this method, one can recover some of the Ornstein-Zernike integral closures, that relate the direct and indirect correlation functions, like the PY closure for instance. Our approach is different from the DFT in that we find a relation between the probability correlation

(density) functions of successive orders, rather than a relation between the direct and indirect correlation functions.

In this paper we have used the maximum entropy principle to derive a closure relation for the BBGKY hierarchy. This approach to the closure problem appears to be new. We proved that for entire space systems, the maximum entropy closure relation coincides with the probabilistic Kirkwood superposition approximation for all orders of the hierarchy. In confined systems, with finite number of particles, the maximum entropy closure differ from the Kirkwood superposition approximation. In particular, when applied to the highest level of the hierarchy, the maximum entropy closure is exact, while the probabilistic Kirkwood approximation is not. Together with the generality of the maximum entropy principle, we are lead to believe that the maximum entropy closure will outperform the Kirkwood superposition approximation, even for low order approximations. We expect to notice the differences between the resulting pair correlation function especially near the domain boundaries. The implementation and empirical results are a subject to a separate paper.

## 10 Acknowledgments

Special thanks to Bob Eisenberg, Douglas Henderson and Roland Roth for reading the manuscript and for their useful comments.

## References

- [1] R. S. Berry, S. Rice, J. Ross, *Physical Chemistry*, 2nd edition. Oxford University Press, NY 2000.
- [2] J.M.G. Barthel, A.H. Baumgartel (Editor), H. Krienke, *Physical Chemistry of Electrolyte Solutions: Modern Aspects*, Steinkopf, Dietrich Pub. 1998.
- [3] H. T. Davis, *Statistical Mechanics of Phases, Interfaces, and Thin Films*, (Advances in Interfacial Engineering Series), John Wiley & Sons, NY, 1995.
- [4] D. Henderson (Editor), *Fundamentals of inhomogeneous fluids*, Marcel Dekker, New York, 1992

- [5] E. L. Lehmann, *Testing Statistical Hypotheses*, 2nd Ed., Springer Verlag, 1997.
- [6] E. L. Lehmann, G. Casella, *Theory of Point Estimation*, 2nd Ed., Springer Verlag, 1998.
- [7] R.S. Eisenberg, "Channels as enzymes." *J. Memb. Biol.* **115**, pp.112 (1990).
- [8] R.S. Eisenberg, "Ion Channels as Devices", *Journal of Computational Electronics* **2**, pp.245-249 (2003).